



Intestinal parasitic infections among children aged 12–59 months in Nyamasheke District, Rwanda



Evariste Hakizimana^{1,2,3,†}, Ju Yeong Kim^{1,†} , Singeun Oh¹, Moonsoo Yoon¹, Tai-Soon Yong^{1,*} 

¹Department of Tropical Medicine, Institute of Tropical Medicine, Arthropods of Medical Importance Resource Bank, Yonsei University College of Medicine, Seoul 03722, Korea; ²Graduate School of Public Health, Yonsei University, Seoul 03722, Korea; ³Kamonyi Health Center, Nyamasheke District, Western Province, Rwanda

Abstract

Received: 10 April 2023

Accepted: 3 May 2023

*Correspondence
(tsyong212@yuhs.ac)

† These authors contributed equally to this work.

Citation

Hakizimana E, Kim JY, Oh S, Yoon M, Yong TS. Intestinal parasitic infections among children aged 12–59 months in Nyamasheke District, Rwanda. Parasites Hosts Dis 2023;61(3):304-309.

Intestinal parasitic infections are a public health burden and a major cause of illness in developing countries. The diseases lead to various health threats, including growth retardation and mental health-related disorders, especially in children. We assessed the risk factors for intestinal parasitic infections among children aged 12–59 months residing in Nyamasheke District, Rwanda. A cross-sectional descriptive study was conducted using secondary data from 1,048 children aged 12–59 months whose stool samples were examined for the presence of intestinal parasites and whose results were registered in the laboratory information system in 2020. The prevalence of intestinal parasites in children aged 12–59 months was 53.2%. The dominant parasites were *Ascaris lumbricoides* (13.1%), followed by *Giardia lamblia* (10.9%), *Entamoeba histolytica* (7.9%), *Trichuris trichiura* (6.5%), hookworms (1.7%), and *Taenia* species (1.4%). A significant association was observed between intestinal parasites and the literacy of mothers or children's caregivers (odds ratio (OR)=5.09, $P < 0.001$). Children from farming households were 2.8-fold more likely to contract intestinal parasitic infections than those from nonfarming households (OR=2.8, $P < 0.001$). A significant association was also observed between intestinal parasites and food safety (OR=4.9, $P < 0.001$). Intestinal parasitic infections were significantly associated with hand hygiene practices after using the toilet and washing fresh fruits before eating ($P < 0.001$). The information gathered will help public health providers and partners develop control plans in highly endemic areas in Rwanda.

Keywords: Intestinal parasite, risk factor, hygiene practice, sanitation

Intestinal parasitic infections are a significant cause of illness and mortality worldwide [1,2]. In developing countries, poor sanitation and sanitary practices increase the risk of contracting these illnesses [3]. Globally, 3.5 billion individuals and 450 million children develop these infections [4]. Children are the most commonly affected age group, and intestinal parasitic infections can affect growth and increase the risk of developing other diseases [5,6]. The common parasites infecting children aged ≤ 5 years from developing countries include *Ascaris lumbricoides*, *Entamoeba histolytica*, and *Giardia lamblia* [7].

Studies in European countries have reported an intestinal parasitic infection prevalence of 5.9% in children [8]. According to the Association of Southeast Asian Nations, 200 million individuals are infected with soil-transmitted helminths, with *A. lumbricoides* being the most common [9-11]. In African countries, including Ethiopia and Nigeria, a significant proportion of children contract intestinal parasitic infections, which have long-term effects on physical growth and cognitive and mental development [12,13]. Intestinal proto-

© 2023 The Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author contributions

Conceptualization: Evariste H, Yong TS
 Data curation: Oh S
 Formal analysis: Kim JY
 Supervision: Yoon M, Yong TS
 Writing: Evariste H, Kim JY

Conflict of interest

The authors have no conflicts of interest.

ORCID

Ju Yeong Kim
 (<https://orcid.org/0000-0003-2456-6298>)
 Tai-Soon Yong
 (<https://orcid.org/0000-0002-3445-0769>)

zoan and helminthic infections are prevalent worldwide; however, their distribution varies based on personal and communal hygiene practices, sanitation, and environmental conditions. The World Health Organization (WHO) reported that more than 95% of parasitic infection cases are because of poverty, low income, lack of awareness, illiteracy, improper hand washing following urination, open defecation practices, contaminated drinking water consumption, improper toilet use, improper food handling, and cultural practices [14].

The WHO has set a regular mass drug administration target for all preschoolers and schoolchildren in 46 targeted African countries, including Rwanda, to reduce the burden of intestinal parasitic infections [15]. Despite these efforts to address intestinal parasitic infections, they remain a significant public health concern in Rwanda. In Nyamasheke District, the rate of medication use for intestinal worms was the highest among children aged 12–59 months, as documented in clinical reports from health facilities in 2020. Therefore, intestinal parasitic infections were the major reason for mothers or children to seek medical attention. However, no scientific study has evaluated the prevalence of intestinal parasitic infections and associated risk factors among children aged 12–59 months in the study area.

In this study, we assessed the prevalence of intestinal parasitic infections and associated factors among children aged 12–59 months in Nyamasheke District, Rwanda. This study provides valuable data to help policymakers, decision makers, and program planners successfully implement interventions that can decrease the public health burden.

This study was conducted in Nyamasheke District, Rwanda. Rwanda is located in Central and East Africa. The Nyamasheke District, with a total area of 1,175 square kilometers, is one of the 7 districts in western Rwanda. A large part of this area is surrounded by Lake Kivu, which is the water source of several households. It is composed of 15 administrative sectors, 20 health centers, and 2 hospitals. The Nyamasheke District has an estimated population of 381,804 residents. However, it is predominantly composed of rural areas, and only a small part is composed of small cities (urban areas). The primary sources of income include agriculture, small businesses, and fishing.

This cross-sectional descriptive study used sociodemographic data, hygiene-related behavior, and stool examination results recorded in a laboratory information system in 2020 during the treatment of children. It was conducted among 1,048 children aged 12–59 months. Their stool samples were examined for the presence of intestinal parasites, and the results were recorded in the laboratory information system for 2020. Using the random multistage sampling method, a total of 1,048 children were selected. The laboratory information system comprises data from 20 health centers and 2 hospitals in this district. Forty-five and seventy-four children were selected from each health center and hospital, respectively.

The sociodemographic characteristics, hygiene-related behavior, and stool examination results of 1,048 children aged 12–59 months recorded in a laboratory information system were analyzed. All results recorded in the database (negative or positive) were considered. Intestinal parasitic infections were used as dependent variables, and their associations with sociodemographics, improper food handling, and poor hand hygiene practices were analyzed. To determine participants' general characteristics, Chi-square test was used. To identify factors associated with intestinal parasitic infections among children aged 12–59 months in this area, binomial logistic regression was used.

Approximately 53.2% of children aged 12–59 months were infected with at least one in-

testinal parasite (Table 1). The most prevalent helminthic parasite was *A. lumbricoides* (13.1%), followed by *T. trichiura* (6.5%), hookworms (1.7%), and *Taenia* species (1.4%) (Table 1). The most prevalent protozoan parasites were *G. lamblia* (10.9%), *E. coli* (10.5%), and *E. histolytica* (7.9%) (Table 1).

The sociodemographic characteristics of children aged 12–59 months with intestinal parasitic infections in Nyamasheke District, Rwanda are presented in Table 2. No significant difference was observed in intestinal parasitic infections by place of residence, sex of children, or children's age group. Of the 1,048 children, 994 lived in rural areas, and 495 (52.4%) were infected with intestinal parasites. Of the 104 children who lived in urban areas, 63 (60.6%) contracted an infection ($P=0.114$). Approximately 52.9% and 53.6% of male and female children contracted infection ($P=0.818$). No significant differences were

Table 1. Prevalence of intestinal parasite infections by species in children aged 12–59 months in Nyamasheke District, Rwanda

Parasite species	No. of infection cases (n=1,048)	Infection cases (%)
<i>Ascaris lumbricoides</i>	137	13.1
<i>Trichuris trichiura</i>	68	6.5
Hookworm	18	1.7
<i>Taenia</i> spp.	15	1.4
<i>Giardia lamblia</i>	114	10.9
<i>Entamoeba histolytica</i>	83	7.9
<i>Entamoeba coli</i>	110	10.5
Any parasite infection	558	53.2

Table 2. Sociodemographic characteristics of children with intestinal parasite infections in Nyamasheke District, Rwanda

Variables		Total number (n=1,048)	Infection cases (%)	P-value ^a
Place of residency	Urban	104	63 (60.6)	0.114
	Rural	944	495 (52.4)	
Sex of children	Male	535	283 (52.9)	0.818
	Female	513	275 (53.6)	
Age group	12–23 months	139	80(57.6)	0.274
	24–59 months	909	478 (52.6)	
Literacy of the mother or child's caretaker	Yes	254	56 (22.0)	<0.001
	No	794	502 (63.2)	
Occupation of the mother or child's caretaker	Not farmer	233	79 (33.9)	<0.001
	Farmer	815	479 (58.8)	
Family size	2–5 people	861	443 (51.5)	0.013
	6–9 people	187	115 (61.5)	
Washing fresh food before cooking	Yes	223	31 (13.9)	<0.001
	No	825	527 (63.9)	
Using piped water and detergent to clean kitchenware	Yes	218	28 (12.8)	<0.001
	No	830	530 (63.9)	
Washing children's hands with soap and water before eating	Yes	168	6 (3.6)	<0.001
	No	880	552 (62.7)	
Washing children's hands with soap and water after using the restroom	Yes	126	8 (6.3)	<0.001
	No	922	550 (59.7)	
Washing fresh fruits before eating	Yes	130	7 (5.4)	<0.001
	No	918	551 (60.0)	

^aChi-square test was performed to calculate the P-value.

observed in intestinal parasitic infections between children aged 12–23 months (57.6%) and those aged 24–59 months (52.6%, $P = 0.274$).

The mothers' literacy and occupation as well as the family size were associated with parasitic infections in children (Table 2). Approximately 63.2% of children with illiterate mothers (or caretakers) were infected with parasites, whereas only 22.0% of children with literate mothers were infected with parasites ($P < 0.001$). Children whose mothers were farmers (58.8%) had more prevalence of parasitic infections than those whose mothers were non-farmers (39%; $P < 0.001$). Larger families (6–9 individuals, 61.5%) had a higher proportion of children with parasitic infections than smaller families (2–5 individuals, 51.5%, $P = 0.013$).

Washing hands, food, and using piped water were not performed by most participants, which caused a significant difference in parasitic infection rates among children (Table 2). Only 13.9% of children (31/223) whose mothers responded that they washed fresh food before cooking were infected with parasites, whereas 63.9% of children (527/825) whose mothers responded that they did not wash the food were infected with parasites ($P < 0.001$). Only 12.8% of children (28/218) who used piped water and detergent to clean their kitchenware were infected with parasites, whereas 63.9% of children (530/830) who did not use piped water to clean kitchenware were infected with parasites, as reported by their mothers ($P < 0.001$). Only 3.6% of children (6/168) whose mothers responded that they washed their hands with soap and water before eating were infected with parasites, whereas 62.7% of children (552/880) whose mothers responded that they did not wash their hands with soap and water were infected with parasites ($P < 0.001$). Only 6.3% of children (8/126) who washed their hands with soap after using the restroom were infected with parasites, whereas 59.7% of children (550/922) who did not were infected with parasites, as reported by their mothers ($P < 0.001$). Only 5.4% of the children (7/130) who washed their fruits before eating were infected with parasites, whereas 60.0% of children (551/918) who did not were infected with parasites, as reported by their mothers ($P < 0.001$).

Multivariate logistic regression was performed using variables that showed significant

Table 3. Multivariate logistic regression analysis of intestinal parasite infections with sociodemographic factors

Variables		Odds ratio	P-value
Literacy of the mother or child's caretaker	Yes	Reference	
	No	5.0995	< 0.001
Occupation of the mother or child's caretaker	Not Farmer	Reference	
	Farmer	2.8519	< 0.001
Family size	2–5 persons	Reference	
	6–9 persons	0.8671	0.515
Washing fresh food before cooking	Washing	Reference	
	Not washing	4.9206	< 0.001
Using pipe water and detergent to clean kitchenware	Yes	Reference	
	No	1.6544	0.126
Washing children's hands with soap and water before eating	Yes	Reference	
	No	8.4507	< 0.001
Washing children's hands with soap and water after using the restroom	Yes	Reference	
	No	4.7458	< 0.001
Washing fresh fruits before eating	Yes	Reference	
	No	9.762	< 0.001

differences (Table 2). The factors associated with the probability of intestinal parasitic infections among children in the Nyamasheke District are shown in Table 3. Children from illiterate families were 5.09-fold more likely to be infected with parasites than those from literate families (odds ratio [OR] = 5.09, $P < 0.001$). The mothers' or child caretakers' occupation (farmer) was associated with intestinal parasitic infections (OR = 2.8, $P < 0.001$). Unwashed fresh food before cooking was associated with intestinal parasitic infections (OR = 4.9, $P < 0.001$). Moreover, failure to wash children's hands with soap and water before eating was associated with intestinal parasitic infections (OR = 8.4, $P < 0.001$). Additionally, failure to wash children's hands with soap after using the toilet was associated with parasitic infections (OR = 4.7, 95% confidence interval: 1.9–11.5). Children from households that did not wash fresh fruits before eating were 9.7-fold more likely to contract parasitic infections than those from households that washed fresh fruits before eating (OR = 9.7, $P < 0.001$). In the multivariate logistic regression model, family size ($P = 0.515$) and using piped water and detergent to clean kitchenware ($P = 0.126$) were not significant factors affecting the risk of parasitic infections (Table 3).

Of the 1,048 children, 558 (53.2%) were infected with intestinal parasites. Chirdan et al. [16] evaluated the prevalence of intestinal parasites in children attending Day Care Centers in Jos, Central Nigeria and reported that 57.8% of children aged 0–59 months were infected with intestinal parasites. In our study, the most prevalent intestinal parasite was *A. lumbricoides* (13.1%); this rate is higher than that reported in a study conducted in a metropolitan city of Bangladesh (1.4% prevalence among pediatric patients) [17] and lower than that reported in a study on ascariasis prevalence conducted among children aged 1–5 years in Markurdi, Benue State, Nigeria (18.75%) [18].

Sociodemographic factors, poor hand hygiene practices, and improper food handling were the primary factors that contributed to the highest intestinal parasitic infection prevalence among children aged 12–59 months in the Nyamasheke District. In our study, the prevalence of intestinal parasitic infections was not significantly different in children from rural and urban areas. However, a study conducted in Kassala Town, Sudan reported that children displaced from rural areas (22.2%) had a higher prevalence of infection than those from urban areas (12.9%) [19]. In the present study, children from households engaged in farming activities had a higher prevalence of intestinal parasitic infections. Similarly, a study conducted in Ethiopia reported that children from families engaged in farming activities had a higher prevalence of intestinal parasitic infections than those from families not engaged in farming activities [20].

To increase the adoption of healthy hygiene practices and lifestyles, health education is necessary. In this study, the education of the mother or child caretaker was associated with intestinal parasitic infections. Intestinal parasitic infections were more prevalent in children from illiterate households.

In conclusion, intestinal parasitic infections remain a public health problem in the Nyamasheke District. Health education campaigns on proper handwashing and food handling should be implemented, particularly in rural areas and among illiterate mothers or child caretakers, to decrease the prevalence of intestinal parasitic infections. To explore other factors associated with intestinal parasitic infections in the Nyamasheke District, further studies are also warranted.

References

- Farrell SH, Coffeng LE, Truscott JE, Werkman M, Toor J, et al. Investigating the effectiveness of current and modified World Health Organization guidelines for the control of soil-transmitted helminth infections. *Clin Infect Dis* 2018;66(suppl):S253-S259. <https://doi.org/10.1093/cid/ciy002>
- Mulatu G, Zeynudin A, Zemene E, Debalke S, Beyene G. Intestinal parasitic infections among children under five years of age presenting with diarrhoeal diseases to two public health facilities in Hawassa, South Ethiopia. *Infect Dis Poverty* 2015; 4:49. <https://doi.org/10.1186/s40249-015-0081-x>
- Campbell SJ, Nery SV, D'Este CA, Gray DJ, McCarthy JS, et al. Water, sanitation and hygiene-related risk factors for soil-transmitted helminth and *Giardia duodenalis* infections in rural communities in Timor-Leste. *Int J Parasitol* 2016;46(12):771-779. <https://doi.org/10.1016/j.ijpara.2016.07.005>
- Arani AS, Alaghebandan R, Akhlaghi L, Shahi M, Lari AR. Prevalence of intestinal parasites in a population in south of Tehran, Iran. *Rev Inst Med Trop Sao Paulo* 2008;50(3):145-149. <https://doi.org/10.1590/s0036-46652008000300003>
- Okay P, Ertug S, Gultekin B, Onen O, Beser E. Intestinal parasites prevalence and related factors in school children, a western city sample-Turkey. *BMC publ Health* 2004;4(1):1-6. <https://doi.org/10.1186/1471-2458-4-64>
- Fauziah N, Aviani JK, Agrianfanny YN, Fatimah SN. Intestinal parasitic infection and nutritional status in children under five years old: A Systematic Review. *Trop Med Infect Dis* 2022; 7(11):371. <https://doi.org/10.3390/tropicalmed7110371>
- El-Sherbini GT, Abosdera MM. Risk factors associated with intestinal parasitic infections among children. *J Egypt Soc Parasitol* 2013;43(1):287-294. <https://doi.org/10.12816/0006385>
- Kantzanou M, Karalexi MA, Vrioni G, Tsakris A. Prevalence of intestinal parasitic infections among children in Europe over the last five years. *Trop Med Infect Dis* 2021;6(3):160. <https://doi.org/10.3390/tropicalmed6030160>
- Bdir S, Adwan G. Prevalence of intestinal parasitic infections in Jenin Governorate, Palestine: a 10-year retrospective study. *Asian Pac J Trop Med* 2010;3(9):745-747. [https://doi.org/10.1016/S1995-7645\(10\)60179-](https://doi.org/10.1016/S1995-7645(10)60179-)
- Khan W, Nisa NU, Khan A. Prevalence and risk factors associated with intestinal parasitic infections among food handlers of Swat, Khyber Pakhtunkhwa, Pakistan. *J Food Nutr Res* 2017; 5:331-336. <https://doi.org/10.12691/jfnr-5-5-7>
- Bakarman MA, Hegazi MA, Butt NS. Prevalence, characteristics, risk factors, and impact of intestinal parasitic infections on school children in Jeddah, Western Saudi Arabia. *J Epidemiol Glob Health* 2019;9(1):81. <https://doi.org/10.2991/jegh.k.190219.001>
- Chelkeba L, Mekonnen Z, Alemu Y, Emanu D. Epidemiology of intestinal parasitic infections in preschool and school-aged Ethiopian children: a systematic review and meta-analysis. *BMC Public Health* 2020;20(1):1-16. <https://doi.org/10.1186/s12889-020-8222-y>
- Karshima SN. Prevalence and distribution of soil-transmitted helminth infections in Nigerian children: a systematic review and meta-analysis. *Infect Dis Poverty* 2018;7(04):1-14. <https://doi.org/10.1186/s40249-018-0451-2>
- Torgerson PR, Devleeschauwer B, Praet N, Speybroeck N, Willingham AL, et al. World Health Organization estimates of the global and regional disease burden of 11 foodborne parasitic diseases, 2010: a data synthesis. *PLoS Med* 2015;12(12): e1001920. <https://doi.org/10.1371/journal.pmed.1001920>
- World Health Organization. Soil-transmitted helminthiasis. Number of children treated 2007-2008: update on the 2010 global target. *Wkly Epidemiol Rec* 2010;85:141-147.
- Chirdan OO, Akosu JT, Adah SO. Intestinal parasites in children attending day care centers in Jos, Central Nigeria. *Niger J Med* 2010;19(2). <https://doi.org/10.4314/njm.v19i2.56526>
- Nipa NJ, Akter N, Hira HM, Akter F, Jahan D, et al. Intestinal parasitic infections among the pediatric patients in a metropolitan city of bangladesh with emphasis on cryptosporidiosis. *Cureus* 2022;14(7):e23316. <https://doi.org/10.7759/cureus.26927>
- Okoh ME, Nyinoh IW, Utume LN, Terzunwe TT. Prevalence of ascariasis among children in makurdi, benue state, nigeria. *J Parasitol Res* 2014;21(5):69-73. <https://doi.org/10.9734/jamb/2021/v21i530353>
- Mohamed MM, Ahmed AI, Salah ET. Frequency of intestinal parasitic infections among displaced children in Kassala Town. *Khartoum Med J* 2012;2(1):175-177.
- Eyayu T, Wubie A, Kiros T, Tiruneh T, Damtie S, et al. Prevalence of intestinal parasitosis and its associated factors among children aged 6 to 59 months attending Mekane Eyesus Primary Hospital, Northcentral Ethiopia. *Glob Pediatr Health* 2021; 8:2333794X211036605. <https://doi.org/10.1177/2333794X211036605>